

R-PEN- penicillin g potassium powder, for solution
Huvepharma, Inc.

R-Pen®

0.5 BU Front Panel

Antibiotic

R-Pen®
(penicillin G potassium)

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

ANTIBIOTIC
for Drinking Water

Nonsterile

Active Ingredient:

0.500 billion units
penicillin G potassium

Mix 1 packet to 333 gallons water

Store between 20°C - 25°C
(68°F - 77°F) with excursions
permitted between 15°C - 40°C
(59°F - 104°F).

For Oral Use in Turkeys Only

Keep Out of Reach of Children

Made in Canada

Huvepharma
Manufactured for:
Huvepharma, Inc.
525 Westpark Dr, Suite 230
Peachtree City, GA 30269

Take Time



Observe Label
Directions

Restricted Drug (California) -

Use only as directed.

Not for human use

Approved by FDA under ANADA # 200-106

KCP-J00003 (Rev. 03-2023)

Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD
104078A

Gravity Flow Watering Systems:

Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

RESIDUE WARNINGS:

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS:

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

INDICATIONS:

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS:

Combine contents of entire package and approximately 1.5 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

Automatic Watering Systems:

Pour the concentrated solution into a glass or plastic container, then add enough water to make 2.6 gallons (9.9 liters) of solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

1.0 BU Front Panel

Antibiotic

R-Pen®

(penicillin G potassium)

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

ANTIBIOTIC for Drinking Water

Nonsterile

Active Ingredient:

1.0 billion units

penicillin G potassium

Mix 1 packet to 666 gallons water

Store between 20°C - 25°C

(68°F - 77°F) with excursions

permitted between 15°C - 40°C

(59°F - 104°F).

For Oral Use in Turkeys Only

Keep Out of Reach of Children

Made in Canada

Huvepharma Manufactured for

Huvepharma, Inc.

525 Westpark Dr, Suite 230

Peachtree City, GA 30269

Take Time



**Observe Label
Directions**

Restricted Drug (California) -

Use only as directed.

Not for human use

Approved by FDA under ANADA # 200-106

KCP-J00004 (Rev. 02-2023)

Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD

104079A

Gravity Flow Watering Systems:

Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

RESIDUE WARNINGS:

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS:

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

INDICATIONS:

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS:

Combine contents of entire package and approximately 3 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

Automatic Watering Systems:

Pour the concentrated solution into a glass or plastic container, then add enough water to make 5.2

gallons (19.7 liters) of solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

0.5 BU Packet Label Image

Gravity Flow Watering Systems:

Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

RESIDUE WARNINGS:

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS:

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

R-Pen[®]
(penicillin G potassium)

**ANTIBIOTIC
for Drinking Water**

Nonsterile

Active Ingredient:
0.500 billion units
penicillin G potassium

Mix 1 packet to 333 gallons water
Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

For Oral Use in Turkeys Only

Keep Out of Reach of Children

Made in Canada



Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

Take Time



**Observe Label
Directions**

Restricted Drug (California).
Use only as directed.
Not for human use.

Approved by FDA under ANADA # 200-106
KCP-J00003 (Rev. 03-2023)

Huvepharma and R-Pen are registered trademarks of
Huvepharma EOOD
104078A

INDICATIONS:

For treatment of erysipelas in turkeys
(caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

DIRECTIONS:

Combine contents of entire package and approximately 1.5 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

Automatic Watering Systems:

Pour the concentrated solution into a glass or plastic container; then add enough water

to make 2.6 gallons (9.9 liters) of solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.



R-Pen[®]

1.0 BU Packet Label Image

NO COPY

Gravity Flow Watering Systems:

Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

RESIDUE WARNINGS:

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

PRECAUTIONS:

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

CORNER RADIUS

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

R-Pen[®]

(penicillin G potassium)

**ANTIBIOTIC
for Drinking Water**

Nonsterile

Active Ingredient:

1.0 billion units
penicillin G potassium

Mix 1 packet to 666 gallons water

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

For Oral Use in Turkeys Only

Keep Out of Reach of Children

Made in Canada



HUVEPHARMA

Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

Take Time



**Observe Label
Directions**

Restricted Drug (California).

Use only as directed.

Not for human use.

Approved by FDA under ANADA # 200-106

KCP-J00004 (Rev. 02-2023)

Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD

104079A

CORNER RADIUS

INDICATIONS:

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

DOSAGE AND ADMINISTRATION:

Administer orally at a dosage of

DIRECTIONS:

Combine contents of entire package and approximately 3 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

add enough water to make 5.2 gallons (19.7 liters) of solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

disappears.

Automatic Watering Systems:
Pour the concentrated solution into a glass or plastic container, then

(50 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.



R-Per[®]

NO COPY

R-PEN

penicillin g potassium powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:23243-6768
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PENICILLIN G POTASSIUM (UNII: VL775ZTH4C) (PENICILLIN G - UNII:Q42T66VG0C)	PENICILLIN G	500000000 [iU] in 500000000 [iU]

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6768-1	25 in 1 PAIL		
1		500000000 [iU] in 1 PACKET		
2	NDC:23243-6768-2	12 in 1 PAIL		
2		10000000000 [iU] in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200106	04/13/2017	

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (552691651)